

## Human SDF1 protein, His tag (active)

**Cat. No. GTX00098-pro****Applications** Functional Assay**Species** Human**Package**

10 µg

## Applications

**Application Note**

SDF1 (stromal cell-derived factor 1), also known as C-X-C motif chemokine 12, is a chemokine protein that has chemotaxis active on T-lymphocytes and monocytes. It is thought that SDF1 stimulates migration of monocytes and T-lymphocytes through its receptors, CXCR4 and ACKR3; thus, chemotaxis assay used 24-well microchemotaxis system was undertaken to detect the chemotactic effect of SDF1 on the human monocytic cell line THP-1. Briefly, THP-1 cells were seeded into the upper chambers (100 µl cell suspension,  $1 \times 10^6$  cells/ml in RPMI-1640 with 0.5% FBS) and SDF1 (10 ng/ml, 30 ng/ml and 60 ng/ml diluted separately in serum free RPMI-1640) was added in lower chamber with a polycarbonate filter (8µm pore size) used to separate the two compartments. After incubation at 37°C with 5% CO<sub>2</sub> for 3h, the filter was removed, then cells in low chamber were observed by inverted microscope at low magnification (×100) and the number of migrated cells were counted at high magnification (×400) randomly (five fields for each filter). The migrated THP-1 cells in low chamber at low magnification (×100) randomly, and the migrated cells were counted at high magnification (×400). SDF1 is able to induce migration of THP-1 cells, and the optimum chemotaxis of SDF1 occurs at 10-60 ng/ml.


**Observed MW (kDa)** 9 kDa.

## Properties

**Form** Lyophilized powder**Buffer** Reconstitute with 20mM Tris and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris, 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose.**Preservative** ProClin 300**Storage** For short-term storage (1-2 weeks), store at 4°C. For long-term storage, store at -20°C or below. After reconstitution, keep as concentrated solution. Avoid freeze-thaw cycles.**Region/Sequence** N-terminal His-Tag; Pro23~Lys89 (NP\_000600.1)**Expression System** E. coli**Purity** > 95%**Endotoxin** < 1 EU/µg**Conjugation** Unconjugated**Note** For laboratory use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

For full product information, images and publications, please visit our [website](#).

DATA IMAGES




**GTX00098-pro Functional Assay Image**

The chemotactic effect of GTX00098-pro Human SDF1 protein (active).

Upper: (A) THP-1 cells were seeded into the upper chambers and serum free RPMI-1640 without SDF1 was added in lower chamber, then cells in lower chamber were observed at low magnification ( $\times 100$ ) after incubation for 3hrs.


(B) THP-1 cells were seeded into the upper chambers and serum free RPMI-1640 with 10 ng/ml SDF1 was added in lower chamber, then cells in lower chamber were observed at low magnification ( $\times 100$ ) after incubation for 3hrs.

Lower: The number of migrated cells were counted at high magnification ( $\times 400$ ) randomly (five fields for each filter).



**GTX00098-pro Image**

SDS-PAGE analysis of GTX00098-pro Human SDF1 protein (active).



**GTX00098-pro Image**

WB analysis of GTX00098-pro Human SDF1 protein (active).



For full product information, images and publications, please visit our [website](#).