

Human EPO protein, His tag (active)

Cat. No. GTX00251-pro**Applications** Functional Assay**Species** Human**Package**

10 µg

Applications

Application Note

Erythropoietin (EPO), also known as hematopoietin or hemopoietin, is a glycoprotein cytokine secreted by the kidney in response to cellular hypoxia. Erythropoietin is an essential hormone for red blood cell production. Its primary effect on red blood cell progenitors and precursors (which are found in the bone marrow in humans) by promoting their survival through protecting these cells from apoptosis, or cell death. EPO is the primary erythropoietic factor that cooperates with various other growth factors involved in the development of erythroid lineage from multipotent progenitors. Besides, Erythropoietin Receptor (EPOR) has been identified as an interactor of EPO, thus a binding ELISA assay was conducted to detect the interaction of recombinant human EPO and recombinant human EPOR. Briefly, EPO were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100 µl were then transferred to EPOR-coated microtiter wells and incubated for 2h at 37°C. Wells were washed with PBST and incubated for 1h with anti-EPO pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37°C. Finally, add 50 µl stop solution to the wells and read at 450nm immediately. The binding activity of EPO and EPOR was in a dose dependent manner.

To test the effect of EPO on cell proliferation, TF-1 cells were seeded into triplicate wells of 96-well plates at a density of 5000 cells/well with 1% serum standard RPMI-1640 including various concentrations of recombinant human EPO. After incubated for 72h, cells were observed by inverted microscope and cell proliferation was measured by Cell Counting Kit-8 (CCK-8). Briefly, 10 µl of CCK-8 solution was added to each well of the plate, then the absorbance at 450nm was measured using a microplate reader after incubating the plate for 1-4 hours at 37°C. Proliferation of TF-1 cells after incubation with EPO for 72h observed by inverted microscope. Cell viability was assessed by CCK-8 (Cell Counting Kit-8) assay after incubation with recombinant EPO for 72h, and EPO significantly increased cell viability of TF-1 cells.

Observed MW (kDa) 33-40 kDa.


Properties

Form Lyophilized powder**Buffer** Reconstitute with 10mM PBS (pH7.4) to 0.1-1.0mg/ml. Do not vortex. Lyophilized from PBS (pH7.4), 0.01% SKL, 1mM DTT, 5% Trehalose.**Preservative** ProClin 300**Storage** For short-term storage (1-2 weeks), store at 4°C. For long-term storage, store at -20°C or below. After reconstitution, keep as concentrated solution. Avoid freeze-thaw cycles.**Region/Sequence** N-terminal His-Tag; Ala28~Arg193 (NP_000790.2)**Expression System** HEK293 cells**Purity** > 95%**Endotoxin** < 1 EU/µg**Conjugation** Unconjugated

For full product information, images and publications, please visit our [website](#).

Note For laboratory use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.


DATA IMAGES



GTX00251-pro Functional Assay Image


Cell proliferation effect of GTX00251-pro Human EPO protein (active). Cell viability was measured by Cell Counting Kit-8 (CCK-8).

- (A) Unstimulated TF-1 cells cultured in RPMI-1640 for 72hrs.
- (B) TF-1 cells cultured in RPMI-1640, stimulated with 50 ng/ml EPO for 72hrs.



GTX00251-pro Functional Assay Image

Functional ELISA analysis of GTX00251-pro Human EPO protein (active) which can bind immobilized EPOR protein.



GTX00251-pro Image

SDS-PAGE analysis of GTX00251-pro Human EPO protein (active).



For full product information, images and publications, please visit our [website](#).