

# Mouse HEXA protein, His tag

# Cat. No. GTX00306-pro

| Applications | Functional Assay | <mark>Package</mark><br>10 μg |
|--------------|------------------|-------------------------------|
| Species      | Mouse            | . 13                          |

### **Applications**

#### **Application Note**

Hexosaminidase A, Alpha (HEXa) is a lysosomal enzyme. There are three predominant isoenzymes: hexosaminidase A, B and S. Hexosaminidase A and the cofactor GM2 activator protein catalyze the degradation of the GM2 gangliosides and other molecules containing terminal N-acetyl hexosamines. The enzymes are composed of two alpha and/or beta subunits, which are coded by HEXA and HEXB genes, respectively. Even though the alpha and beta subunits of hexosaminidase A can both cleave GalNAc residues, only the alpha subunit which contains a key residue, Arg-424 is able to hydrolyze GM2 gangliosides. Hexosaminidase A (alpha polypeptide) plays a critical role in the brain and spinal cord (central nervous system). Besides, Hexosaminidase B Beta (HEXB) has been identified as an interactor of HEXA, thus a binding ELISA assay was conducted to detect the interaction of recombinant mouse HEXA and recombinant mouse HEXB. Briefly, HEXA were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100 µl were then transferred to HEXB-coated microtiter wells and incubated for 2h at 37°C. Wells were washed with PBST and incubated for 1h with anti-HEXA pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37°C. Finally, add 50 µl stop solution to the wells and read at 450nm immediately. The binding activity of of HEXA and HEXB was in a dose dependent manner.

Observed MW (kDa) 28 kDa.

| Properties        |  |  |
|-------------------|--|--|
| Form              | Lyophilized powder   |  |
| Buffer            | Reconstitute with 20mM Tris and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris, 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose.                   |  |
| Preservative      | ProClin 300  |  |
| Storage           | For short-term storage (1-2 weeks), store at 4°C. For long-term storage, store at -20°C or below. After reconstitution, keep as concentrated solution. Avoid freeze-thaw cycles. |  |
| Region/Sequence   | N-terminal His-Tag; Leu319~Thr528 (NP_034551.2)  |  |
| Expression System | E. coli  |  |
| Purity            | > 97%  |  |
| Endotoxin         | < 1 EU/μg  |  |
| Conjugation       | Unconjugated   |  |
| Note              | For laboratory use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.   |  |




For full product information, images and publications, please visit our website.

Date 2026 / 01 / 02 Page 1 of 2




# DATA IMAGES




### **GTX00306-pro Functional Assay Image**

Functional ELISA analysis of GTX00306-pro Mouse HEXA protein which can bind immobilized HEXB protein.



### GTX00306-pro Image

SDS-PAGE analysis of GTX00306-pro Mouse HEXA protein.



# GTX00306-pro Image

WB analysis of GTX00306-pro Mouse HEXA protein.



For full product information, images and publications, please visit our <u>website</u>.

Date 2026 / 01 / 02 Page 2 of 2