

Rat ODC protein, His tag

Cat. No. GTX00369-pro

Applications Functional Assay**Species** Rat**Package**

10 µg

Applications

Application Note


The enzyme ornithine decarboxylase (ODC) catalyzes the decarboxylation of ornithine (a product of the urea cycle) to form putrescine. This reaction is the committed step in polyamine synthesis. Lysine 69 on ornithine decarboxylase (ODC) binds the cofactor pyridoxal phosphate to form a Schiff base. Ornithine displaces the lysine to form a Schiff base attached to ODC, which decarboxylates to form a quinoid intermediate. This intermediate rearranges to form a Schiff base attached to putrescine, which is attacked by lysine to release putrescine product and reform PLP-bound ODC. Besides, Thymidine Kinase 1, Soluble (TK1) has been identified as an interactor of ODC, thus a binding ELISA assay was conducted to detect the interaction of recombinant rat ODC and recombinant rat TK1. Briefly, ODC were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100 µl were then transferred to TK1-coated microtiter wells and incubated for 2h at 37°C. Wells were washed with PBST and incubated for 1h with anti-ODC pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37°C. Finally, add 50 µl stop solution to the wells and read at 450nm immediately. The binding activity of ODC and TK1 was in a dose dependent manner.

Observed MW (kDa) 55 kDa.


Properties

Form Lyophilized powder**Buffer** Reconstitute with 20mM Tris and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris, 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose.**Preservative** ProClin 300**Storage** For short-term storage (1-2 weeks), store at 4°C. For long-term storage, store at -20°C or below. After reconstitution, keep as concentrated solution. Avoid freeze-thaw cycles.**Region/Sequence** Full length protein, N-terminal His-Tag; Met1~Val461 (NP_001289012.1)**Expression System** E. coli**Purity** > 97%**Endotoxin** < 1 EU/µg**Conjugation** Unconjugated**Note** For laboratory use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.


For full product information, images and publications, please visit our [website](#).

DATA IMAGES

GTX00369-pro Functional Assay Image

Functional ELISA analysis of GTX00369-pro Rat ODC protein which can bind immobilized TK1 protein.


GTX00369-pro Image

WB analysis of GTX00369-pro Rat ODC protein.


GTX00369-pro Image

SDS-PAGE analysis of GTX00369-pro Rat ODC protein.



For full product information, images and publications, please visit our [website](https://www.genetex.com).