

CD11b antibody [ICRF44]

Cat. No. GTX00481

Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Applications	WB, ICC/IF, IHC-Fr, FCM, IP, Activation, Neutralizing /Inhibition
Reactivity	Human, Pig, Baboon, Chimpanzee, Cynomolgus monkey, Rhesus Monkey

References (12)

Package

100 µg

PRODUCT

Summary

The ICRF44 antibody reacts with human CD11b, also known as integrin α M. This 165-170 kDa cell surface glycoprotein is part of a family of integrin α receptors that mediate adhesion between cells (cell-cell) and components of the extracellular matrix, e.g. fibrinogen (cell-matrix). In addition, integrin α s are active signaling receptors which recruit leukocytes to inflammatory sites and promote cell activation. Complete, functional integrin α receptors consist of distinct combinations of integrin α chains which are differentially expressed. integrin α M (CD11b) assembles with integrin α beta-2 (CD18) into a receptor known as Macrophage Antigen-1 (Mac-1) or complement receptor type 3 (CR3). This receptor binds and induces intracellular signaling through ICAM-1, ICAM-2, ICAM-3 and ICAM-4 on endothelial cells and can also facilitate removal of iC3b bearing foreign cells.

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	Assay dependent
ICC/IF	Assay dependent
IHC-Fr	Assay dependent
FCM	Assay dependent
IP	Assay dependent
Activation	Assay dependent
Neutralizing /Inhibition	Assay dependent

Not tested in other applications.

Calculated MW 127 kDa. ([Note](#))

Properties

Form	Liquid
Buffer	10mM NaH ₂ PO ₄ , 150mM NaCl
Preservative	0.09% Sodium azide



For full product information, images and publications, please visit our [website](#).

Date 2026 / 05 / 04 Page 1 of 2

Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C.
Concentration	0.5 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Rheumatoid synovial cells and human monocytes.
Purification	Purified by affinity chromatography From tissue culture supernatant
Conjugation	Unconjugated
Note	<p>For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.</p> <p>Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.</p>



For full product information, images and publications, please visit our [website](#).