

CD45R antibody [RA3-6B2]

Cat. No. GTX00486

Host	Rat	
Clonality	Monoclonal	
Isotype	lgG2a	
Applications	WB, ICC/IF, IHC-P, IHC-Fr, FCM, IP, Depletion	
Reactivity	Human, Mouse, Cat	

References (8) Package 100 µg

PRODUCT

Summary

The RA3-6B2 antibody reacts with the human and mouse CD45 isoform known as CD45R, or B220, a protein tyrosine phosphatase of \geq 220 kDa. CD45 is one of the most abundant hematopoietic markers, and is expressed on all leukocytes (the Leukocyte Common Antigen, LCA). Various isoforms are generated and expressed in cell-specific patterns, all critical for leukocyte function. In mouse, the CD45R/B220 isoform is predominantly found on B cells, at varying levels on all stages from pro-B cells to activated B cells, and may also be detected on certain T cell and NK cell subsets. It is of note that B220 is not similarly expressed on human B cells, where it appears to be differentiation-specific and therefore expressed on only some B cell subsets. Other forms of CD45 with restricted cellular expression include CD45RA, CD45RB, CD45RO and several others.

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution	
WB	Assay dependent	
ICC/IF	Assay dependent	
IHC-P	Assay dependent	
IHC-Fr	Assay dependent	
FCM	Assay dependent	
IP	Assay dependent	
Depletion	Assay dependent	
Not tested in other applications.		

Calculated MW 145 kDa. (Note)

Product Note

This antibody recognizes an epitope on CD45R, which is expressed at all developmental stages of B cells, including activated B cells, but also on subsets of NK and T cells. T cells detected by this antibody are supposed to be in activated state.

P	'ro	pe	rties

Form Liquid



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 10 / 28 Page 1 of 2



Buffer	10mM NaH ₂ PO ₄ , 150mM NaCl
Preservative	0.09% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C.
Concentration	0.5 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Abelson murine leukemia virus-induced pre-B tumor cells
Purification	Purified by affinity chromatography From tissue culture supernatant
Conjugation	Unconjugated
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our website.

Date 2025 / 10 / 28 Page 2 of 2