

NUP98 antibody [13C2]

Cat. No. GTX00693

Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Applications	WB, ICC/IF
Reactivity	Human, Saccharomyces cerevisiae, Schizosaccharomyces pombe, Tetrahymena

References (1)

Package

100 µg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	Assay dependent
ICC/IF	Assay dependent

Not tested in other applications.

Product Note

For ICC/IF application, it works on *S. cerevisiae*, *S. pombe*, and *T. etrahymena* but is not suitable for human. Recommend GTX00695 NUP98 antibody [21A10] on human samples.

Properties

Form	Liquid
Buffer	Filter-sterilized PBS, 50% Glycerol
Preservative	No preservatives
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Synthetic peptides containing conserved N-terminal sequence, GLFG, of Nup98 protein of Tetrahymena thermophila: Peptide 1; 1-MFGNTGGGGLFGNTQTQQTGGGLFGQPQQ-29 Peptide 2; 646-SNPTQGGGLFGAANPGLGG-664 Epitope determined: FGxxN
Purification	Purified IgG
Conjugation	Unconjugated




For full product information, images and publications, please visit our [website](#).

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Note

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

DATA IMAGES



GTX00693 ICC/IF Image

ICC/IF analysis of *S. cerevisiae* cells using GTX00693 NUP98 antibody [13C2], GTX00695 NUP98 antibody [21A10], or GTX00697 NUP98 antibody [2H10].

Green : Primary antibody

Violet : DAPI

Dilution :


13C2 or 21A10 : 1:10

2H10 : 10 µg/ml

SUMMARY OF THE SUITABILITY OF MABS FOR IMMUNOLOGICAL APPLICATIONS									
mAb	Isotype	<i>T. thermophilus</i>		HeLa cell		<i>S. cerevisiae</i>		<i>S. pombe</i>	
		IF	WB	IF	WB	IF	WB	IF	WB
13C2	Mouse IgG1	+++	+++	+	+++	+++	+++	+++	+++
21A10	Mouse IgG1	+++	+	+++	-	+++	++	+++	++
2H10	Rat IgG2c	-	+	+++	+	+++	+++	+++	++
									Unidentified

GTX00693 Image

Summary of the suitability of GTX00693 NUP98 antibody [13C2], GTX00695 NUP98 antibody [21A10], or GTX00697 NUP98 antibody [2H10] for immunological applications.



GTX00693 ICC/IF Image


ICC/IF analysis of HeLa cells using GTX00693 NUP98 antibody [13C2], GTX00695 NUP98 antibody [21A10], or GTX00697 NUP98 antibody [2H10]. The signal at the nuclear periphery with 21A10 mAb was much higher and the background lower than that of 2H10 and 13C2 antibodies.

Green : Primary antibody

Violet : DAPI

Dilution : 0.5 µg/ml

Fixation : Cold Methanol (-30 degree C) for 30 min



GTX00693 ICC/IF Image

ICC/IF analysis of *S. pombe* cells using GTX00693 NUP98 antibody [13C2], GTX00695 NUP98 antibody [21A10], or GTX00697 NUP98 antibody [2H10].

Green : Primary antibody

Violet : DAPI

Fixation : 4% PFA for 10 min, treated with 0.6 mg/ml Zymolyase 100T at 3 degree C for 70 min

Permeabilization : 1% Triton X-100 for 1 min



For full product information, images and publications, please visit our [website](https://www.genetex.com).