

Histone H4K5ac (acetyl Lys5) antibody [GT1138]

Cat. No. GTX00775

Host	Rabbit
Clonality	Monoclonal
Isotype	IgG
Application	WB, ICC/IF, IHC-P
Reactivity	Human, Mouse, Rat

Package
100 µl

APPLICATION

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500 - 1:2000
ICC/IF	1:50 - 1:200
IHC-P	1:50 - 1:200

Not tested in other applications.

PROPERTIES

Form	Liquid
Buffer	PBS, 50% Glycerol
Preservative	0.02% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	Batch dependent (Please refer to the vial label for the specific concentration.)
Immunogen	A synthesized peptide derived from human Histone H4 (acetyl K5)
Purification	Purified by affinity chromatography
Conjugation	Unconjugated

Note


For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.




For full product information, images and publications, please visit our [website](#).

DATA IMAGES



GTX00775 WB Image

Untreated (–) and treated (+) HeLa whole cell extracts (30 μg) were separated by 15% SDS-PAGE, and the membrane was blotted with Histone H4K5ac (acetyl Lys5) antibody [GT1138] (GTX00775) diluted at 1:500. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.




GTX00775 WB Image

WB analysis of HeLa and C6 whole cell lysates using GTX00775 Histone H4K5ac (acetyl Lys5) antibody [GT1138].

Dilution : 1:1000

Loading : 25μg




GTX00775 ICC/IF Image

ICC/IF analysis of C6 cells treated with/without TSA (1 μM at 37°C for 18 hours) using GTX00775 Histone H4K5ac (acetyl Lys5) antibody [GT1138].

Orange: Primary antibody

Blue: DAPI



GTX00775 ICC/IF Image

ICC/IF analysis of U2OS cells treated with/without TSA (1 μM at 37°C for 18 hours) using GTX00775 Histone H4K5ac (acetyl Lys5) antibody [GT1138].

Orange: Primary antibody

Blue: DAPI



For full product information, images and publications, please visit our [website](https://www.genetex.com).