

GAK antibody [9-13]

Cat. No. GTX00873

Host	Mouse
Clonality	Monoclonal
Isotype	lgG1
Application	WB, ICC/IF, IHC-P
Reactivity	Human, Rat

Package 100 μΙ

APPLICATION

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500
ICC/IF	1:50
IHC-P	Assay dependent
Not tested in other applications.	

Calculated MW 143 kDa. (<u>Note</u>)

PROPERTIES	
Form	Liquid
Buffer	PBS, 50% Glycerol
Preservative	No preservative
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	Batch dependent (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant rat GAK (kinase domain, 1-430)
Purification	Purified IgG From culture supernant
Conjugation	Unconjugated
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.




For full product information, images and publications, please visit our website.

Date 2024 / 04 / 29 Page 1 of 2




DATA IMAGES



GTX00873 IHC-P Image

IHC-P analysis of human prostate cancer tissue section using GTX00873 GAK antibody [9-13]. It showed nuclear accumulation of GAK in cancer cells.


Dilution: 1:50



GTX00873 ICC/IF Image

ICC/IF analysis of TIG-1 cells using GTX00873 GAK antibody [9-13].

Dilution: 1:50



GTX00873 WB Image

WB analysis of TIG-1 and PC-3 whole cell lysates using GTX00873 GAK antibody [9-13].

Dilution: 1:500



For full product information, images and publications, please visit our website.

Date 2024 / 04 / 29 Page 2 of 2