

GAK antibody [9-10]

Cat. No. GTX00874

Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Applications	WB, ICC/IF, IHC-P
Reactivity	Human, Rat

Package
100 µl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500
ICC/IF	1:50
IHC-P	Assay dependent

Not tested in other applications.

Calculated MW 143 kDa. ([Note](#))

Properties

Form	Liquid
Buffer	PBS, 50% Glycerol
Preservative	No preservatives
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	Batch dependent (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant rat GAK (kinase domain, 1-430)
Purification	Purified IgG From culture supernant
Conjugation	Unconjugated

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Note


Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).


Date 2026 / 01 / 07 Page 1 of 2

DATA IMAGES

**GTX00874 ICC/IF Image**


ICC/IF analysis of TIG-1 cells using GTX00874 GAK antibody [9-10].

Dilution : 1:50

**GTX00874 WB Image**

WB analysis of TIG-1, HeLa S3 and PC-3 whole cell lysates using GTX00874 GAK antibody [9-10].

Dilution : 1:500

**GTX00874 IHC-P Image**

IHC-P analysis of human prostate cancer tissue section using GTX00874 GAK antibody [9-10]. It shows nuclear accumulation of GAK in cancer cells.

Dilution : 1:50



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 07 Page 2 of 2