

## CDC6 antibody

## Cat. No. GTX00896

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, ICC/IF, IP, ChIP assay
Reactivity	Human, Mouse, Rat

## Package

100 µg

## Applications

## Application Note

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:1000-1:3000
ICC/IF	1:200
IP	1:200
ChIP assay	Assay dependent

Not tested in other applications.

Calculated MW 63 kDa. ([Note](#))

## Properties

Form	Liquid
Buffer	Filter-sterilized PBS, 50% Glycerol
Preservative	No preservatives
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant GST-human CDC6 (1-326 amino acids) expressed in E. coli
Purification	Purified by antigen-affinity chromatography From serum
Conjugation	Unconjugated




For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 09 Page 1 of 2

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

**Note**

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.


**DATA IMAGES****GTX00896 IP Image**

IP analysis of HeLa whole cell lysate using GTX00896 CDC6 antibody.

Lane 1 : Control IP with non-immune IgG

Lane 2 : Immunoprecipitates with anti-CDC6 antibody


Lane 3 : Input

**GTX00896 WB Image**

WB analysis of HeLa whole cell lysate using GTX00896 CDC6 antibody.

Dilution : 1:1000

Loading : 10 µg

**GTX00896 ICC/IF Image**

ICC/IF analysis of MEF cells (E14.5) using GTX00896 CDC6 antibody. Note that CDC6 abundantly localizes in nuclei of the cells at G1 and early S phase.

Fixation : 4% PFA, RT, 10 min

Permeabilization : ice-cold methanol, on ice, 10 min

Dilution : 1:200



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 09 Page 2 of 2