

LC3B antibody [GT1187]

Cat. No. GTX00949

Host	Rabbit
Clonality	Monoclonal
Isotype	IgG
Applications	WB, ICC/IF, IHC-P
Reactivity	Human, Mouse, Rat

References (2)

Package

100 µl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500 - 1:2000
ICC/IF	1:50 - 1:200
IHC-P	1:50 - 1:200

Not tested in other applications.

Calculated MW	15 kDa. (Note)
Observed MW (kDa)	14, 16 kDa. The observed M.W. is based on the following publication. PMID: 31576176

Properties

Form	Liquid
Buffer	PBS, 0.05% BSA, 50% Glycerol
Preservative	0.05% ProClin 300
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	Batch dependent (Please refer to the vial label for the specific concentration.)
Immunogen	A synthetic peptide of human LC3B.
Purification	Purified by affinity chromatography
Conjugation	Unconjugated




For full product information, images and publications, please visit our [website](#).

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Note


Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

DATA IMAGES



GTx00949 WB Image

Untreated (–) and treated (+) HepG2 whole cell extracts (30 μg) were separated by 15% SDS-PAGE, and the membrane was blotted with LC3B antibody [GT1187] (GTx00949) diluted at 1:500. The HRP-conjugated anti-rabbit IgG antibody (GTx213110-01) was used to detect the primary antibody.



GTx00949 ICC/IF Image

ICC/IF analysis of PFA-fixed HeLa cells with/without Chloroquine (50 μM, 37°C, 20 hrs) treatment using GTx00949 LC3B antibody [GT1187].


Right : untreated HeLa cells

Left : HeLa cells with 50 μM Chloroquine (20 hrs, 37°C) treatment

Orange : Primary antibody

Blue : DAPI

Dilution : 1:100



GTx00949 ICC/IF Image

ICC/IF analysis of PFA-fixed NIH-3T3 cells with/without Chloroquine (50 μM, 37°C, 20 hrs) treatment using GTx00949 LC3B antibody [GT1187].

Right : untreated NIH-3T3 cells

Left : NIH-3T3 cells with 50 μM Chloroquine (20 hrs, 37°C) treatment

Orange : Primary antibody

Blue : DAPI

Dilution : 1:100



For full product information, images and publications, please visit our [website](https://www.genetex.com).