

IL17A antibody

Cat. No. GTX01094

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, ICC/IF, IHC-P, ELISA
Reactivity	Human, Mouse

Package 100 μg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	0.125-0.5 μg/mL
ICC/IF	5-20 μg/mL
IHC-P	2-5 μg/mL
ELISA	Assay dependent

Not tested in other applications.

Calculated MW 18 kDa. (Note)

Properties	
Form	Liquid
Buffer	PBS
Preservative	0.02% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Peptide corresponding to 16 amino acids near the center of mature human IL-17. The immunogen is located within amino acids 50-100 of IL-17.
Purification	Purified by antigen-affinity chromatography
Conjugation	Unconjugated



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 12 / 16 Page 1 of 2




For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Note

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

DATA IMAGES




GTX01094 WB Image

WB analysis of IL-17 recombinant protein using GTX01094 IL17A antibody.

Loading: 30 ng of human IL-17 recombinant protein per lane

Dilution : 0.125 μ g/mL (Lane 1) ; 0.25 μ g/mL (Lane 2) ; 0.5 μ g/mL (Lane 3)




GTX01094 IHC-P Image

IHC-P analysis of formaldehyde-fixed human tonsil tissue using GTX01094 IL17A antibody.

Antigen retrieval: Heat mediation with a citrate buffer (pH6)

Dilution: $5\mu g/mL$



GTX01094 IHC-P Image

ICC/IF analysis of 4% PFA-fixed A-20 cells using GTX01094 IL17A antibody.

Green: Primary antibody

Blue : DAPI

Dilution: 5 μg/mL



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 12 / 16 Page 2 of 2