

ASIC3 antibody

Cat. No. GTX01177

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, FCM
Reactivity	Human, Mouse, Rat

Package
100 µg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	0.1-0.5µg/ml
FCM	1-3µg/1x10 ⁶ cells

Not tested in other applications.

Calculated MW 59 kDa. ([Note](#))

Product Note Superfamily members of ASIC3 are not reactive to this antibody.

Properties

Form	Liquid
Buffer	5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄
Preservative	0.05mg Thimerosal, 0.05mg Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.5 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	A synthetic peptide corresponding to a sequence at the N-terminus of human ASIC3(56-73aa FLYQVAERVRYYREFHHQ).
Purification	Purified by antigen-affinity chromatography
Conjugation	Unconjugated

Note


For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

DATA IMAGES



GTX01177 FCM Image


FACS analysis of U937 cells using GTX01177 ASIC3 antibody.

Blue : Primary antibody

Green : Isotype control

Red : No antibody control

Antibody amount : 1µg/1x10⁶ cells



GTX01177 WB Image

WB analysis of various sample lysates using GTX01177 ASIC3 antibody.

Lane 1 : Rat Brain Tissue Lysate

Lane 2 : Rat Testis Tissue Lysate

Lane 3 : U87 Cell Lysate

Lane 4 : NEURO Cell Lysate

Lane 5 : SMMC Cell Lysate

Loading : 50µg of sample under reducing conditions

Dilution : 0.5 µg/mL



For full product information, images and publications, please visit our [website](#).