

Integrin alpha 2 antibody [HMa2] (APC)

Cat. No. GTX01474-07

Host Armenian Hamster**Clonality** Monoclonal**Isotype** IgG**Applications** FCM**Reactivity** Mouse

References (4)

Package

100 µg

PRODUCT

Summary

The HMa2 antibody reacts with mouse CD49b, a 150 kDa glycoprotein member of the integrin of family. CD49b, also known as integrin α 2, associates non-covalently with CD29 (integrin β 1) to form the VLA-2 heterodimer. VLA-2 integrin α function as cellular receptors for collagen and laminin. CD49b is expressed by platelets, intestinal epithelial cells, NK cells, epithelial cells, and a subset of CD4+ T cells. CD49b plays an essential role in mediating the adhesion of cells to the extracellular matrix.

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution

Recommended dilution

FCM

Assay dependent

Not tested in other applications.

Properties

Form Liquid**Buffer** 10mM NaH₂PO₄, 150mM NaCl, 0.1% Gelatin**Preservative** 0.09% Sodium azide**Storage** Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C. DO NOT FREEZE. Protect from light.**Concentration** 0.2 mg/ml (Please refer to the vial label for the specific concentration.)**Purification** Purified by affinity chromatography
From tissue culture supernatant**Conjugation** Allophycocyanin (APC) [Wavelength](#)

Note


For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

DATA IMAGES

**GTX01474-07 FCM Image**

FACS analysis of mouse C57Bl/6 splenocytes using GTX01474-07 Integrin alpha 2 antibody [HMa2] (APC).

Right panel : co-stained with Integrin alpha 2 antibody [HMa2] (APC) and Mouse NK1.1 antibody (FITC)

Left panel : co-stained with isotype control and Mouse NK1.1 antibody (FITC)

antibody amount : 0.125 μ g (5 μ l)



For full product information, images and publications, please visit our [website](#).