

# CD40 antibody [FGK4.5/ FGK45] (PE)

### Cat. No. GTX01483-08

| Host         | Rat        |
|--------------|------------|
| Clonality    | Monoclonal |
| Isotype      | lgG2a      |
| Applications | FCM        |
| Reactivity   | Mouse      |

References (4)
Package
100 µg

#### PRODUCT

# Summary

The FGK45 antibody reacts with mouse CD40, a 40-50 kDa glycoprotein member of the tumor necrosis factor receptor (TNFR) family. CD40 is constitutively expressed by antigen presenting cells, including dendritic cells, B cells, and macrophages. The interaction of CD40 and its ligand CD154, expressed on activated T cells, plays an essential role in mediating a broad range of immune and inflammatory responses including T cell dependent immunoglobulin class switching, germinal center formation, and the development of memory B cells. The FGK45 clone is reported to be functionally useful for both in vitro and in vivo activation of CD40 expressing antigen presenting cells. The FGK45 antibody reacts with mouse CD40, a 40-50 kDa glycoprotein member of the tumor necrosis factor receptor (TNFR) family.

#### **Applications**

#### **Application Note**

\*Optimal dilutions/concentrations should be determined by the researcher.

| Suggested dilution | Recommended dilution |
|--------------------|----------------------|
| FCM                | Assay dependent      |

Not tested in other applications.

| Properties    |   |
|---------------|---|
| Form          | Liquid  |
| Buffer        | 10mM NaH <sub>2</sub> PO <sub>4</sub> , 150mM NaCl, 0.1% Gelatin  |
| Preservative  | 0.09% Sodium azide  |
| Storage       | Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C. DO NOT FREEZE. Protect from light.  |
| Concentration | 0.2 mg/ml (Please refer to the vial label for the specific concentration.)  |
| Purification  | Purified by affinity chromatography From tissue culture supernatant   |
| Conjugation   | Phycoerythrin (PE) Wavelength   |
| Note          | For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.                           |
|               | Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product. |




For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 07 / 12 Page 1 of 2



## DATA IMAGES



#### GTX01483-08 FCM Image

FACS analysis of mouse C57Bl/6 splenocytes using GTX01483-08 CD40 antibody [FGK45] (PE).

Solid lone : primary antibody Dashed line : isotype control antibody amount : 0.25  $\mu$ g (5  $\mu$ l)



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 07 / 12 Page 2 of 2