## CD40L / CD154 antibody [5C8]

## Cat. No. GTX01484

Host	Mouse	References (5)
Clonality	Monoclonal	<mark>Package</mark> 100 μg
lsotype	lgG2a	
Applications	FCM	
Reactivity	Human, Cynomolgus monkey, Rhesus Monkey	

PRODUCT	
Summary	The 5C8 antibody reacts with human CD154 which is also known as CD40L (CD40 ligand). CD154 is a 39kDa member of the TNF-α superfamily and is predominantly expressed on activated T cells but is also expressed on NK cells, mast cells, basophils, and eosinophils. CD154 binds to CD40, a molecule that is expressed on antigen presenting cells including B cells, monocytes/macrophages, and dendritic cells. The interaction of CD154 and CD40 regulates the activation of antigen presenting cells and is critical for B cell costimulation. The 5C8 antibody has been reported to block CD40L-CD40 interactions.
Applications	

## **Application Note**

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
FCM	Assay dependent

Not tested in other applications.

Properties	
Form	Liquid
Buffer	10mM NaH <sub>2</sub> PO <sub>4</sub> , 150mM NaCl
Preservative	0.09% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C.
Concentration	0.5 mg/ml (Please refer to the vial label for the specific concentration.)
Purification	Purified by affinity chromatography From tissue culture supernatant
Conjugation	Unconjugated
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our <u>website</u>.