

CD80 antibody [2D10.4]

Cat. No. GTX01487

Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Applications	ICC/IF, FCM, Activation
Reactivity	Human, Rhesus Monkey

References (3)

Package

100 µg

PRODUCT

Summary

The 2D10.4 antibody reacts with human CD80, a 60 kDa glycoprotein that is a member of the Ig superfamily. CD80, also known as B7-1, is a ligand for the T cell receptors CD28 and CTLA-4 (CD152). Along with CD86, CD80 plays a major role in regulating the activation of T cells. CD80 is expressed by activated antigen presenting cells including B cells, macrophages, and dendritic cells. Activated T cells have also been reported to express CD80.

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
ICC/IF	Assay dependent
FCM	Assay dependent
Activation	Assay dependent

Not tested in other applications.

Properties

Form	Liquid
Buffer	10mM NaH ₂ PO ₄ , 150mM NaCl
Preservative	0.09% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C.
Concentration	0.5 mg/ml (Please refer to the vial label for the specific concentration.)
Purification	Purified by affinity chromatography From tissue culture supernatant
Conjugation	Unconjugated



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 31 Page 1 of 2

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Note

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 31 Page 2 of 2