

PD-L1 antibody [29E.2A3] (APC)

Cat. No. GTX01495-07

Host	Mouse
Clonality	Monoclonal
Isotype	IgG2b
Applications	FCM
Reactivity	Human

References (6)

Package

100 test

PRODUCT

Summary

The 29E.2A3 antibody reacts with human CD274 which is also known as PD-L1 or B7-H1. CD274 is a type 1 transmembrane glycoprotein and a member of the Ig superfamily. PD-L1 is a ligand for CD279 (PD-1) and plays an important role in regulating immune cell responses. CD274 is expressed on activated monocytes, macrophages, dendritic cells, and T cells.

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution

Recommended dilution

FCM

2 µg (5 µl) for 10⁵-10⁸ cells in 100 µl sample per test

Not tested in other applications.

Properties

Form Liquid

Buffer 10mM NaH₂PO₄, 150mM NaCl, 0.1% Gelatin

Preservative 0.09% Sodium azide

Storage Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C. DO NOT FREEZE. Protect from light.

Concentration 0.4 mg/ml (Please refer to the vial label for the specific concentration.)

Purification Purified by affinity chromatography
From tissue culture supernatant

Conjugation Allophycocyanin (APC) [Wavelength](#)

Note


For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

DATA IMAGES

**GTX01495-07 FCM Image**

FACS analysis of human PBMCs (PHA stimulation for 3 days) using GTX01495-07 PD-L1 antibody [29E.2A3] (APC).

Solid line : primary antibody

Dashed line : isotype control

antibody amount : 2 μ g (5 μ l)



For full product information, images and publications, please visit our [website](#).