

Cyclin D1 antibody [ZM178]

Cat. No. GTX01660

Host	Mouse	
Clonality	Monoclonal	
Isotype	lgG2a	
Application	IHC-P, Neutralizing/Inhibition	
Reactivity	Human	

Package 500 μl

APPLICATION

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution	
IHC-P	1:100-1:200	
Neutralizing/Inhibition	Assay dependent	
Nets - Destructions of descriptions of discounties have induced orders and since a serious law assessment in the serious law assessment of the serious law as the serious law assessment of the serious law as the serious		

Note: Pretreatment of deparaffinized tissue with heat-induced epitope retrieval or enzymatic retrieval is recommended. This antibody neutralizes the activity of cyclin D1 in vivo.

Not tested in other applications.

Calculated MW 34 kDa. (Note)

PROPERTIES	
Form	Liquid
Buffer	Tris-HCl buffer, 0.2% BSA
Preservative	0.09% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C.
Concentration	Batch dependent (Please refer to the vial label for the specific concentration.)
Immunogen	Synthetic peptide corresponding to residues within aa 200-295 of human Cyclin D1
Purification	Purified IgG
Conjugation	Unconjugated
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our <u>website</u>.

Date 2024 / 04 / 27 Page 1 of 1