

## Cyclin D1 antibody [ZM178]

**Cat. No. GTX01660**

<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Isotype</b>	IgG2a
<b>Applications</b>	IHC-P, Neutralizing /Inhibition
<b>Reactivity</b>	Human

**Package**  
500 µl

## Applications

**Application Note**

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
IHC-P	1:100-1:200
Neutralizing /Inhibition	Assay dependent

**Note : Pretreatment of deparaffinized tissue with heat-induced epitope retrieval or enzymatic retrieval is recommended. This antibody neutralizes the activity of cyclin D1 in vivo.**

Not tested in other applications.

## Properties

<b>Form</b>	Liquid
<b>Buffer</b>	Tris-HCl, 0.2% BSA
<b>Preservative</b>	0.09% Sodium azide
<b>Storage</b>	Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C.
<b>Concentration</b>	Batch dependent (Please refer to the vial label for the specific concentration.)
<b>Immunogen</b>	Synthetic peptide corresponding to residues within aa 200-295 of human Cyclin D1
<b>Purification</b>	Purified IgG
<b>Conjugation</b>	Unconjugated

**Note**

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).