

Myotilin antibody [RSO34]

Cat. No. GTX01881

Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Applications	IHC-P, IHC-Fr
Reactivity	Human

Package
500 µl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
IHC-P	1:40–1:80
IHC-Fr	1:20

Note : High temperature antigen unmasking technique. Standard ABC technique. Standard ABC technique. Acetone fixation recommended.

Not tested in other applications.

Properties

Form	Liquid
Buffer	Tissue culture supernatant
Preservative	0.09% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Immunogen	Prokaryotic recombinant protein corresponding to a C-terminal region of 266 amino acids of the human myotilin molecule.
Purification	Unpurified
Conjugation	Unconjugated

Note


For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

DATA IMAGES

**GTX01881 IHC-P Image**

IHC-P analysis of human skeletal muscle tissue using GTX01881 Myotilin antibody [RSO34]. Note sarcoplasmic staining of normal muscle fibers (A) and presence of protein aggregates (B).



For full product information, images and publications, please visit our [website](#).