

NCAM antibody [CD564]

Cat. No. GTX01917

Host	Mouse
Clonality	Monoclonal
Isotype	IgG2b
Applications	IHC-P
Reactivity	Human

Package
500 µl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
IHC-P	1:100

Note : The recommended fixative is 10% neutral-buffered formalin for paraffin-embedded tissue sections. Heat Induced Epitope Retrieval (HIER) was performed in Epitope Retrieval Solution pH6.

Not tested in other applications.

Properties


Form	Liquid
Buffer	Tissue culture supernatant
Preservative	0.09% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C. DO NOT FREEZE.
Immunogen	Prokaryotic recombinant protein corresponding to a region of the extracellular domain of the human CD56 molecule.
Purification	Unpurified
Conjugation	Unconjugated
For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.	
Note	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 13 Page 1 of 2

DATA IMAGES

**GTx01917 IHC-P Image**

IHC-P analysis of human tonsil tissue using GTx01917 NCAM antibody [CD564]. Note the NK cells and CD4/CD8 double positive T cells show a weak to moderate and distinct membrane staining reaction while the majority of lymphocytes are unstained.



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 13 Page 2 of 2