

Bcl-2 antibody [bcl-2/100/D5]

Cat. No. GTX01935

Host	Mouse
Clonality	Monoclonal
Isotype	lgG1
Applications	IHC-P
Reactivity	Human

Package 500 μl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
IHC-P	1:100

Note: The recommended fixative is 10% neutral-buffered formalin for paraffin-embedded tissue sections. Heat Induced Epitope Retreival (HIER) was performed in Epitope Retreival Solution pH6.

Not tested in other applications.

Properties	
Form	Liquid
Buffer	Tissue culture supernatant
Preservative	0.09% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C. DO NOT FREEZE.
Immunogen	Synthetic peptide sequence (GAAPAPGIFSSQPGC-COOH).
Purification	Unpurified
Conjugation	Unconjugated
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 11 / 07 Page 1 of 2



DATA IMAGES



GTX01935 IHC-P Image

IHC-P analysis of human follicular lymphoma using GTX01935 Bcl-2 antibody [bcl-2/100/D5]. Neoplastic cells show a moderate, predominantly cytoplasmic staining reaction.



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 11 / 07 Page 2 of 2

€ 886-3-6208988 🔒 886-3-6208989 🐷 infoasia@genetex.com