

# BCL6 antibody [LN22]

**Cat. No. GTX01937**

<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Isotype</b>	IgG2b
<b>Applications</b>	IHC-P
<b>Reactivity</b>	Human

**Package**  
500 µl

## Applications

### Application Note

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
IHC-P	1:60

**Note :** Heat Induced Epitope Retrieval (HIER) was performed in Epitope Retrieval Solution pH 9. The use of H<sub>2</sub>O<sub>2</sub> to block endogenous peroxidase has a detrimental effect on the epitope recognized by this antibody. Solutions containing 1% or greater H<sub>2</sub>O<sub>2</sub> can dramatically reduce the level of staining achievable. Blocking after unmasking can also lead to negative results. Therefore, it is recommended that endogenous peroxidase should be blocked before unmasking with 0.5% H<sub>2</sub>O<sub>2</sub>/methanol for 10 minutes. If this recommendation is not followed, staining intensity may be reduced.

Not tested in other applications.

## Properties

<b>Form</b>	Liquid
<b>Buffer</b>	Tissue culture supernatant
<b>Preservative</b>	0.09% Sodium azide
<b>Storage</b>	Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C. DO NOT FREEZE.
<b>Immunogen</b>	Prokaryotic recombinant protein corresponding to 1–350 amino acids of the N-terminus of the Bcl-6 human oncoprotein molecule.
<b>Purification</b>	Unpurified
<b>Conjugation</b>	Unconjugated

### Note


For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

## DATA IMAGES



### GTx01937 IHC-P Image

IHC-P analysis of human diffuse large B cell lymphoma using GTx01937 BCL6 antibody [LN22]. The majority of neoplastic cells show a moderate to strong nuclear staining reaction.



For full product information, images and publications, please visit our [website](#).