

## Perforin antibody [5B10]

## Cat. No. GTX01949

Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Applications	IHC-P
Reactivity	Human

Package  
500 µl

## Applications

## Application Note

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
IHC-P	1:20

**Note : The recommended fixative is 10% neutral-buffered formalin for paraffin-embedded tissue sections. Heat Induced Epitope Retrieval (HIER) was performed in Epitope Retrieval Solution pH6.**

Not tested in other applications.

**Product Note** This antibody shows cross-reactivity with smooth muscle and differentiated epithelium.

## Properties


Form	Liquid
Buffer	Tissue culture supernatant
Preservative	0.09% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C. DO NOT FREEZE.
Immunogen	Recombinant prokaryotic protein corresponding to an external region of the C-terminus of the perforin molecule.
Purification	Unpurified
Conjugation	Unconjugated
For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.	
Note	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 14 Page 1 of 2

## DATA IMAGES

**GTx01949 IHC-P Image**

IHC-P analysis of human follicular lymphoma using GTx01949 Perforin antibody [5B10]. Note focal granular staining of occasional cytotoxic T lymphocytes.



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 14 Page 2 of 2