

SPTBN2 antibody [SPTBN2/2887R]

Cat. No. GTX02717

Host	Rabbit
Clonality	Monoclonal
Isotype	IgG
Applications	WB, ICC/IF, IHC-P, FCM
Reactivity	Human

Package
100 µg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1-2 µg/ml
ICC/IF	1-2 µg/ml
IHC-P	1-2 µg/ml
FCM	1-2 µg/million cells

Note : Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris buffer with 1mM EDTA (pH 9.0) for 45 min at 95°C followed by cooling at RT for 20 minutes.

Not tested in other applications.

Calculated MW 271 kDa. ([Note](#))

Properties

Form	Liquid
Buffer	PBS, 0.05% BSA
Preservative	0.05% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C.
Concentration	200 µg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant human SPTBN2 fragment (aa356-475)
Purification	Protein A/G purified
Conjugation	Unconjugated




For full product information, images and publications, please visit our [website](#).

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Note

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.


DATA IMAGES

**GTX02717 ICC/IF Image**

ICC/IF analysis of HeLa cells using GTX02717 SPTBN2 antibody [SPTBN2/2887R].

Green : Primary antibody


Red : nuclear counterstain

**GTX02717 FCM Image**

FACS analysis of HeLa cells using GTX02717 SPTBN2 antibody [SPTBN2/2887R].

Blue : Primary antibody

Red : Isotype control

**GTX02717 WB Image**

WB analysis of PANC-1 whole cell lysate using GTX02717 SPTBN2 antibody [SPTBN2/2887R].



For full product information, images and publications, please visit our [website](#).