

c-Myc antibody [GT1265]

Cat. No. GTX02877

Host	Rabbit
Clonality	Monoclonal
Isotype	IgG
Applications	WB, ICC/IF
Reactivity	Human, Mouse

Package
100 µl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	Assay dependent
ICC/IF	Assay dependent

Not tested in other applications.

Calculated MW 49 kDa. ([Note](#))

Properties

Form	Liquid
Buffer	PBS, 0.05% BSA, 50% Glycerol
Preservative	0.02% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	Batch dependent (Please refer to the vial label for the specific concentration.)
Immunogen	A synthesized peptide derived from human c-Myc
Purification	Purified by affinity chromatography
Conjugation	Unconjugated

Note


For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).


DATA IMAGES

**GTX02877 WB Image**

WB analysis of mouse ovary tissue lysate using GTX02877 c-Myc antibody [GT1265].

Dilution : 1:1000

Loading : 25µg per lane


**GTX02877 ICC/IF Image**

ICC/IF analysis of HeLa cells using GTX02877 c-Myc antibody [GT1265].

Red : Primary antibody

Blue : DAPI

Dilution : 1:100

**GTX02877 WB Image**

WB analysis of various cell lysates using GTX02877 c-Myc antibody [GT1265].

Dilution : 1:1000

Loading : 25µg per lane



For full product information, images and publications, please visit our [website](#).