

CD16a antibody

Cat. No. GTX03379

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	ICC/IF, IHC-P, FCM, ELISA
Reactivity	Human

Package 100 μg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
ICC/IF	Assay dependent
IHC-P	Assay dependent
FCM	Assay dependent
ELISA	Assay dependent

Not tested in other applications.

Properties	
Form	Liquid
Buffer	4mg Trehalose, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄
Preservative	0.05mg Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	.5 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	E. coli-derived human CD16 recombinant protein (Position: Q101-D166).
Purification	Purified IgG
Conjugation	Unconjugated
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.




For full product information, images and publications, please visit our website.

Date 2025 / 12 / 12 Page 1 of 2



DATA IMAGES




GTX03379 IHC-P Image

IHC-P analysis of human rectal cancer tissue using GTX03379 CD16a antibody.

Antigen retireval: Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution).

Dilution: 2 μg/ml




GTX03379 ICC/IF Image

ICC/IF analysis of K562 cells using GTX03379 CD16a antibody.

Antigen retireval: IHC enzyme antigen retrieval reagent for 15min.

Dilution : $5 \mu g/mL$




GTX03379 IHC-P Image

IHC-P analysis of human liver cancer tissue using GTX03379 CD16a antibody.

Antigen retireval : Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval

solution).

Dilution : $2 \, \mu g/ml$



GTX03379 FCM Image

FACS analysis of human PBMC cells using GTX03379 CD16a antibody.

Blue: Primary antibody Green: Isotype control> Red: Unlabelled sample

Antibody amount : $1 \mu g/1x10^6$ cells



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 12 / 12 Page 2 of 2