

Lysozyme antibody

Cat. No. GTX03467

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, ICC/IF, IHC-P
Reactivity	Human, Mouse, Rat

Package 100 μg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	0.1-0.5μg/ml
ICC/IF	5μg/ml
IHC-P	0.5-1μg/ml
Note: Antigen retireval by heat	

Not tested in other applications.

Calculated MW 17 kDa. (<u>Note</u>)

Properties	
Form	Liquid
Buffer	5mg BSA, 0.9mg NaCl, 0.2mg Na₂HPO₄
Preservative	0.05mg sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.5 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human Lysozyme (106-141aa; NIADAVACAKRVVRDPQGIRAWVAWRNRCQNRDVRQ).
Purification	Purified by antigen-affinity chromatography
Conjugation	Unconjugated



For full product information, images and publications, please visit our website.

Date 2025 / 12 / 27 Page 1 of 2




For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Note

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

DATA IMAGES




GTX03467 IHC-P Image

IHC-P analysis of human colon organoid tissue using GTX03467 Lysozyme antibody.

Antigen retireval: Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins.

Blue : DAPI Dilution : 5μg/ml




GTX03467 IHC-P Image

IHC-P analysis of human ileum tissue using GTX03467 Lysozyme antibody.

Antigen retireval: Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins.

Blue : DAPI Dilution : 5µg/ml



GTX03467 IHC-P Image

IHC-P analysis of human intestinal cancer tissue using GTX03467 Lysozyme antibody.

Antigen retireval: Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval

solution).

Dilution: 1 μg/ml



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 12 / 27 Page 2 of 2