

# CD93 antibody

**Cat. No. GTX03672**

<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	IgG
<b>Applications</b>	IHC-P, FCM
<b>Reactivity</b>	Human

**Package**  
100 µg

## Applications

### Application Note

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
IHC-P	1-2 µg/ml
FCM	1-3 µg/1x10 <sup>6</sup> cells

Not tested in other applications.

## Properties

<b>Form</b>	Liquid
<b>Buffer</b>	4mg Trehalose, 0.9mg NaCl, 0.2mg Na <sub>2</sub> HPO <sub>4</sub>
<b>Preservative</b>	No preservatives
<b>Storage</b>	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
<b>Concentration</b>	0.5 mg/ml (Please refer to the vial label for the specific concentration.)
<b>Immunogen</b>	A synthetic peptide corresponding to a sequence of human CD93 (QNAADSYSWVPERAESRAMENQ).
<b>Purification</b>	Purified by antigen-affinity chromatography
<b>Conjugation</b>	Unconjugated

### Note


For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

## DATA IMAGES




### GTX03672 IHC-P Image

IHC-P analysis of human colonic adenocarcinoma tissue using GTX03672 CD93 antibody.

Antigen retrieval : Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution)

Dilution : 2 µg/ml




### GTX03672 IHC-P Image

IHC-P analysis of human breast cancer tissue using GTX03672 CD93 antibody.

Antigen retrieval : Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution)

Dilution : 2 µg/ml




### GTX03672 IHC-P Image

IHC-P analysis of human placenta tissue using GTX03672 CD93 antibody.

Antigen retrieval : Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution)

Dilution : 2 µg/ml




### GTX03672 IHC-P Image

IHC-P analysis of human appendiceal adenocarcinoma tissue using GTX03672 CD93 antibody.

Antigen retrieval : Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution)

Dilution : 2 µg/ml



### GTX03672 IHC-P Image

IHC-P analysis of human gall bladder adenosquamous carcinoma tissue using GTX03672 CD93 antibody.

Antigen retrieval : Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution)

Dilution : 2 µg/ml



For full product information, images and publications, please visit our [website](#).