

Goat Anti-Human IgM (Mu chain) antibody, pre-adsorbed (TRITC)

Cat. No. GTX04120-25

Host	Goat
Clonality	Polyclonal
Isotype	IgG
Applications	WB, ICC/IF, IHC-P, IHC-Fr, FCM, IP, ELISA, Activation, Depletion, ELISPOT, Multiplexing, Purification
Reactivity	Human

Package

1 mg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	Assay dependent
ICC/IF	Assay dependent
IHC-P	Assay dependent
IHC-Fr	Assay dependent
FCM	Assay dependent
IP	Assay dependent
ELISA	1:100-1:400
Activation	Assay dependent
Depletion	Assay dependent
ELISPOT	Assay dependent
Multiplexing	Assay dependent
Purification	Assay dependent

Not tested in other applications.

Product Note

Pre-adsorbed with Human IgG and IgA. May react with IgM from other species.

Properties


Form	Liquid
Buffer	PBS
Preservative	0.09% sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C. Protect from light.



For full product information, images and publications, please visit our [website](#).

Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Pooled antisera from goats hyperimmunized with human IgM
Purification	Purified by antigen-affinity chromatography
Conjugation	Tetramethylrhodamine (TRITC) Wavelength
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption. Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

DATA IMAGES



GTX04120-25 ELISA Image

ELISA analysis of purified human immunoglobulins using serially diluted GTX04120-25 Goat Anti-Human IgM (Mu chain) antibody, pre-adsorbed (TRITC).



For full product information, images and publications, please visit our [website](#).