

# Goat Anti-Human IgA (Heavy chain) antibody, F(ab')2 fragment, pre-adsorbed (FITC)

## Cat. No. GTX04127-06

Host	Goat
Clonality	Polyclonal
Isotype	IgG F(ab')2
Applications	IHC-Fr, FCM, ELISA, Activation, ELISPOT
Reactivity	Human

### Package

500 µg

## Applications

### Application Note

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
IHC-Fr	Assay dependent
FCM	≤ 1 µg/10 <sup>6</sup> cells
ELISA	1:100-1:400
Activation	Assay dependent
ELISPOT	Assay dependent

**Note : The suggested use of these reagents is in a final volume of 100µl.**

Not tested in other applications.

**Product Note** Pre-adsorbed with Human IgG and IgM. May react with IgA from other species.

## Properties

Form	Liquid
Buffer	PBS
Preservative	0.09% sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C. Protect from light.
Concentration	0.5 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Pepsin digest of Goat Anti-Human IgA.
Purification	Purified IgG
Conjugation	Fluorescein isothiocyanate (FITC) <a href="#">Wavelength</a>




For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 28 Page 1 of 2

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

**Note**

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

**DATA IMAGES****GTX04127-06 ELISA Image**

ELISA analysis of purified human immunoglobulins using serially diluted GTX04127-06 Goat Anti-Human IgA (Heavy chain) antibody, F(ab')2 fragment, pre-adsorbed (FITC).



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 28 Page 2 of 2