

Goat Anti-Human lambda light chain antibody, F(ab')₂ fragment, pre-adsorbed (FITC)

Cat. No. GTX04135-06

Host	Goat
Clonality	Polyclonal
Isotype	IgG F(ab') ₂
Applications	WB, FCM, ELISA, Activation
Reactivity	Human

Package

500 µg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	Assay dependent
FCM	≤ 1 µg/10 ⁶ cells
ELISA	1:200-1:400
Activation	Assay dependent

Note : The suggested use of these reagents is in a final volume of 100µl.

Not tested in other applications.

Product Note Pre-adsorbed with Human κ light chains. May react with λ light chains from other species.

Properties

Form	Liquid
Buffer	PBS
Preservative	0.09% sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C. Protect from light.
Concentration	0.5 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Pepsin digest of Goat Anti-Human Lambda.
Purification	Purified IgG
Conjugation	Fluorescein isothiocyanate (FITC) Wavelength




For full product information, images and publications, please visit our [website](#).

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Note
 Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

DATA IMAGES



GTX04135-06 ELISA Image

ELISA analysis of purified human immunoglobulins using serially diluted GTX04135-06 Goat Anti-Human lambda light chain antibody, F(ab')2 fragment, pre-adsorbed (FITC).



For full product information, images and publications, please visit our [website](#).