

Goat Anti-Mouse IgM (Mu chain) antibody, F(ab')2 fragment, pre-adsorbed

Cat. No. GTX04194

| | |
|--------------|---|
| Host | Goat |
| Clonality | Polyclonal |
| Isotype | IgG F(ab')2 |
| Applications | WB, ICC/IF, IHC-P, IHC-Fr, FCM, ELISA, Activation, Purification |
| Reactivity | Mouse |

Package
500 µg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

| Suggested dilution | Recommended dilution |
|--------------------|----------------------|
| WB | Assay dependent |
| ICC/IF | Assay dependent |
| IHC-P | Assay dependent |
| IHC-Fr | Assay dependent |
| FCM | Assay dependent |
| ELISA | Assay dependent |
| Activation | Assay dependent |
| Purification | Assay dependent |

Not tested in other applications.

Product Note Pre-adsorbed with Mouse IgG and IgA; human immunoglobulins and pooled sera. May react with immunoglobulins from other species.

Properties

| | |
|---------------|---|
| Form | Liquid |
| Buffer | Borate buffered saline |
| Preservative | No preservatives |
| Storage | Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C. |
| Concentration | 0.5 mg/ml (Please refer to the vial label for the specific concentration.) |
| Immunogen | Pepsin digest of Goat Anti-Mouse IgM. |
| Purification | Purified IgG |



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 29 Page 1 of 2

Conjugation

Unconjugated

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Note

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 29 Page 2 of 2