

Goat Anti-Mouse IgG (Fc) antibody, pre-adsorbed (Biotin)

Cat. No. GTX04197-02

Host	Goat
Clonality	Polyclonal
Isotype	IgG
Applications	WB, ELISA
Reactivity	Mouse

Package
1 mg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	Assay dependent
ELISA	1:5000-1:20000

Not tested in other applications.

Product Note

Reacts with the Fc region of mouse IgG1, IgG2a, IgG2b, IgG2c, and IgG3. Pre-adsorbed with Mouse IgG Fab, IgM, and IgA. May react with immunoglobulins from other species

Properties

Form	Liquid
Buffer	PBS
Preservative	0.09% sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C.
Concentration	0.5 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Pooled antisera from goats hyperimmunized with mouse IgG
Purification	Purified by antigen-affinity chromatography
Conjugation	Biotin

Note


For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

DATA IMAGES

**GTX04197-02 ELISA Image**

ELISA analysis of purified mouse immunoglobulins using serially diluted GTX04197-02 Goat Anti-Mouse IgG (Fc) antibody, pre-adsorbed (Biotin) followed by Streptavidin (HRP).



For full product information, images and publications, please visit our [website](#).