

## Hemoglobin beta S antibody [23E5.H6.G6.C1.H7.F7.G9.F6]

**Cat. No. GTX04268**

<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Isotype</b>	IgG3
<b>Applications</b>	WB, FCM, ELISA, Lateral Flow
<b>Reactivity</b>	Human

**Package**  
100 µg

## Applications

**Application Note**

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1 µg/mL
FCM	Assay dependent
ELISA	1:20000
Lateral Flow	Assay dependent

Not tested in other applications.

<b>Calculated MW</b>	16 kDa. ( <a href="#">Note</a> )
----------------------	----------------------------------

**Product Note** This antibody detect the E6V mutant in the hemoglobin beta subunit and does not react with the HbA, HbF, HbC, or HbA-2 isoform.

## Properties

<b>Form</b>	Liquid
<b>Buffer</b>	20mM Potassium Phosphate, 150mM NaCl
<b>Preservative</b>	0.01% sodium azide
<b>Storage</b>	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
<b>Concentration</b>	1 mg/ml (Please refer to the vial label for the specific concentration.)
<b>Immunogen</b>	Synthetic peptide corresponding to amino acid residues near the N-terminus of Hb beta-subunit conjugated to KLH.
<b>Purification</b>	Protein A purified From tissue culture supernatant
<b>Conjugation</b>	Unconjugated




For full product information, images and publications, please visit our [website](#).

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

**Note**  
Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.


DATA IMAGES



**GTX04268 WB Image**

WB analysis of various sample lysates using GTX04268 Hemoglobin beta S antibody [23E5.H6.G6.C1.H7.F7.G9.F6].


- Lane 1 : Prestained molecular weight marker
  - Lane 2 : HbA
  - Lane 3 : HbA2
  - Lane 4 : HbC
  - Lane 5 : HbF
  - Lane 6 : HbS
  - Lane 7 : BSA
- Dilution : 1:1000  
Loading : 10µg



**GTX04268 WB Image**

WB analysis of various sample lysates using GTX04268 Hemoglobin beta S antibody [23E5.H6.G6.C1.H7.F7.G9.F6].

- Lane 1 : Prestained molecular weight marker
  - Lane 2 : HbA peptide conjugated to BSA
  - Lane 3 : HbA-2 peptide conjugated to BSA
  - Lane 4 : HbC peptide conjugated to BSA
  - Lane 5 : HbF peptide conjugated to BSA
  - Lane 6 : HbS peptide conjugated to BSA
  - Lane 7 : BSA alone
- Loading : 50ng per lane  
Dilution : 1µg/mL



**GTX04268 Lateral Flow Image**

Lateral flow analysis of recombinant HbA (left group) or HbS (right group) proteins using GTX04267 Hemoglobin beta antibody [14G2.G11.F11] and GTX04268 Hemoglobin beta S antibody [23E5.H6.G6.C1.H7.F7.G9.F6].

Antibody amount : GTX04267 Hemoglobin beta antibody 0.5µL at 250ug/mL and GTX04268 Hemoglobin beta S antibody 0.5µL at 1mg/mL



For full product information, images and publications, please visit our [website](#).