

PCID2 antibody

Cat. No. GTX04896

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, IHC-P
Reactivity	Human, Mouse

Package
100 µl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:200-1:1000
IHC-P	1:20-1:100

Not tested in other applications.

Calculated MW 46 kDa. ([Note](#))

Properties

Form	Liquid
Buffer	PBS, 40% Glycerol
Preservative	0.05% Sodium Azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	2.4 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Fusion protein of human PCID2
Purification	Purified by antigen-affinity chromatography
Conjugation	Unconjugated


Note

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.




For full product information, images and publications, please visit our [website](#).

DATA IMAGES

GTx04896 WB Image

WB analysis of TM4 whole cell lysate using GTx04896 PCID2 antibody.


Loading : 40µg

Dilution : 1:250


GTx04896 IHC-P Image

IHC-P analysis of human gastric cancer tissue using GTx04896 PCID2 antibody.

Dilution : 1:25


GTx04896 IHC-P Image

IHC-P analysis of human breast cancer tissue using GTx04896 PCID2 antibody.

Dilution : 1:25



For full product information, images and publications, please visit our [website](#).