

Prostein antibody [MSVA-460R] HistoMAX™

Cat. No. GTX04965

Host	Rabbit
Clonality	Monoclonal
Isotype	IgG
Applications	IHC-P
Reactivity	Human

References (1)

Package

500 µl, 100 µl

PRODUCT

Summary

[Go to Normal Tissue Gallery](#)[Go to Cancer Tissue Gallery](#)

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution

Recommended dilution

IHC-P

1:100 - 1:200

Note : Heat-induced antigen retrieval for 5 minutes in an autoclave at 121°C in pH 7.8 Tris-EDTA-based Target Retrieval Solution buffer.

Not tested in other applications.

Product Note

Highly recommended for IHC-P in human tissues.

Properties

Form	Liquid
Buffer	10mM PBS, BSA (Please contact us for sodium azide-free or sodium azide and BSA-free format)
Preservative	sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	66.66 µg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Synthesized peptides to the N-terminus of human SLC45A3 protein
Purification	Protein A/G purified
Conjugation	Unconjugated

For full product information, images and publications, please visit our [website](#).


Date 2026 / 10 / 08 Page 1 of 2

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Note

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.


DATA IMAGES



GTx04965 IHC-P Image

IHC-P analysis of human gastric mucosa tissue using GTx04965 Prostein antibody [MSVA-460R] HistoMAX™.


Stomach mucosa with granular cytoplasmic protein staining of surface epithelial cells.



GTx04965 IHC-P Image

IHC-P analysis of human prostatic adenocarcinoma (PAC) tissue (Gleason 3+3=6) using GTx04965 Prostein antibody [MSVA-460R] HistoMAX™.

Prostatic adenocarcinoma (Gleason 3+3=6) with strong granular perinuclear cytoplasmic protein staining of tumor cells.



GTx04965 IHC-P Image

IHC-P analysis of human prostate tissue using GTx04965 Prostein antibody [MSVA-460R] HistoMAX™.

Normal prostate with strong granular perinuclear cytoplasmic protein staining of luminal cells.



For full product information, images and publications, please visit our [website](#).