

MOLT-4 whole cell lysate

Cat. No. GTX05056

Species Human

Package

500 µg

PRODUCT

Summary

Cells were washed with PBS and then incubated on ice in modified RIPA buffer to lyse the cells. Protein integrity was ensured using a cocktail of protease inhibitors with broad specificity for the inhibition of aspartic, cysteine, and serine proteases as well as aminopeptidases (0.1 mM AEBSF HCl, 0.08 µM Aprotinin, 5 µM Bestatin, 1.5 µM E-64, 2 µM Leupeptin Hemisulfate, 1 µM Pepstatin A). Phosphatase inhibitors 1 mM NaF and 1 mM Na₃VO₄ were also added. Cell debris was removed by centrifugation. Protein concentration was determined by a modified Lowry assay using a commercially available kit. Protein concentration was adjusted to 4 mg/ml with 1X RIPA buffer. For separation by SDS-PAGE and subsequent western blot analysis, lysates should be diluted by user to desired concentration in SDS-PAGE buffer with 2-mercaptoethanol or dithiothreitol as the reducing agent and heated to 95° C for 5 minutes.

Properties

Form Liquid

Buffer 1X RIPA Buffer with HALT Protease and Phosphatase Inhibitors

Preservative No preservatives

Storage This whole cell lysate is stable for three months at -20°C


Concentration 4 mg/ml (Please refer to the vial label for the specific concentration.)

Note

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

DATA IMAGES



GTX05056 Image

SDS-PAGE analysis of GTX05056 MOLT-4 whole cell lysate.

Lane 1 : GTX05056

Lane 2 : Prestained molecular marker

Lane 3 : Non-reduced MOLT-4 whole cell lysate

Loading : 10 µg



For full product information, images and publications, please visit our [website](#).