

## BAK antibody [N1N2], N-term

## Cat. No. GTX100063

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, IHC-P
Reactivity	Human

References ( 9 )

 Review ( 1 )

Package

100 µl, 25 µl

## Applications

## Application Note

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500-1:3000
IHC-P	1:100-1:1000

Not tested in other applications.

Calculated MW 23 kDa. ( [Note](#) )

## Properties


Form	Liquid
Buffer	PBS, 20% Glycerol
Preservative	0.01% Thimerosal
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Carrier-protein conjugated synthetic peptide encompassing a sequence within the N-terminus region of human BAK. The exact sequence is proprietary.
Purification	Purified by antigen-affinity chromatography.
Conjugation	Unconjugated
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).


Date 2026 / 01 / 09 Page 1 of 2

## DATA IMAGES



## GTX100063 WB Image


Wild-type (WT) and BAK knockout (KO) CHO cell extracts (30 µg) were separated by 12% SDS-PAGE, and the membrane was blotted with BAK antibody [N1N2], N-term (GTX100063) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



## GTX100063 IHC-P Image


BAK antibody [N1N2], N-term detects BAK protein at mitochondria by immunohistochemical analysis.  
Sample: Paraffin-embedded human colon.

BAK stained by BAK antibody [N1N2], N-term (GTX100063) diluted at 1:500.  
Antigen Retrieval: Citrate buffer, pH 6.0, 15 min



## GTX100063 WB Image

Various whole cell extracts (30 µg) were separated by 12% SDS-PAGE, and the membrane was blotted with BAK antibody [N1N2], N-term (GTX100063) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody, and the signal was developed with Trident ECL plus-Enhanced.



## GTX100063 IHC-P Image

Immunohistochemical analysis of paraffin-embedded human liver, bile duct, using BAK1(GTX100063) antibody(10 µg/ml).  
Antigen Retrieval: Trilogy™ (EDTA based, pH 8.0) buffer, 15min



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 09 Page 2 of 2