

## GAD67 antibody

**Cat. No. GTX101881**

<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	IgG
<b>Applications</b>	WB, ICC/IF, IHC-P, IHC-Fr, IHC-Wm, IHC, IHC (Free Floating)
<b>Reactivity</b>	Human, Mouse, Rat, Zebrafish, Monkey

References ( 12 )

Package

100 µl, 25 µl

## Applications

**Application Note**

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500-1:3000
ICC/IF	1:100-1:1000
IHC-P	1:100-1:1000
IHC-Fr	1:100-1:1000
IHC-Wm	Assay dependent
IHC	Assay dependent
IHC (Free Floating)	Assay dependent

Not tested in other applications.

**Calculated MW** 67 kDa. ( [Note](#) )**Product Note** This antibody is specific for GAD67 protein, and it does not cross-react with GAD65 protein.

## Properties

<b>Form</b>	Liquid
<b>Buffer</b>	PBS, 1% BSA, 20% Glycerol
<b>Preservative</b>	0.025% ProClin 300
<b>Storage</b>	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
<b>Concentration</b>	0.22 mg/ml (Please refer to the vial label for the specific concentration.)
<b>Immunogen</b>	Recombinant protein encompassing a sequence within the N-terminus region of human GAD67. The exact sequence is proprietary.
<b>Purification</b>	Purified by antigen-affinity chromatography.



For full product information, images and publications, please visit our [website](#).

**Conjugation**


Unconjugated

**Note**


For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

## DATA IMAGES

**GTX101881 WB Image**

Various tissue extracts (30 µg) were separated by 7.5% SDS-PAGE, and the membrane was blotted with GAD67 antibody (GTX101881) diluted at 1:500. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.


**GTX101881 IHC Image**

GAD67 antibody detects GAD67 protein on myofiber membranes by immunohistochemical analysis.

Sample: Agarose-embedded zebrafish embryo

GAD67 antibody (GTX101881) dilution: 1:200.

Image provided with permission courtesy of Dr. T. Schilling at UC, Irvine.

**GTX101881 IHC-Wm Image**

GAD67 antibody detects GAD67 protein on myofiber membranes by immunohistochemical analysis.

Sample: Whole-mount zebrafish embryo

GAD67 antibody (GTX101881) dilution: 1:200.



For full product information, images and publications, please visit our [website](https://www.genetex.com).