

BAG2 antibody

Cat. No. GTX102406

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Application	WB, IHC-P
Reactivity	Human

Package 100 μl, 25 μl

APPLICATION

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500-1:3000
IHC-P	1:100-1:1000

Not tested in other applications.

Calculated MW 24 kDa. (Note)

PROPERTIES		
Form	Liquid	
Buffer	PBS, 20% Glycerol	
Preservative	0.01% Thimerosal	
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.	
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)	
Immunogen	Recombinant protein encompassing a sequence within the center region of human BAG2. The exact sequence is proprietary.	
Purification	Purified by antigen-affinity chromatography.	
Conjugation	Unconjugated	
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.	
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.	




For full product information, images and publications, please visit our <u>website</u>.

Date 2024 / 04 / 29 Page 1 of 2




DATA IMAGES




GTX102406 WB Image

Wild-type (WT) and BAG2 knockout (KO) HeLa cell extracts (30 µg) were separated by 12% SDS-PAGE, and the membrane was blotted with BAG2 antibody (GTX102406) diluted at 1:1000. The HRP-conjugated antirabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



GTX102406 WB Image

Sample (30 ug of whole cell lysate) A: A431 B: HeLa 12% SDS PAGE GTX102406 diluted at 1:1000



GTX102406 IHC-P Image

Immunohistochemical analysis of paraffin-embedded U87 xenograft, using BAG2(GTX102406) antibody at 1:500 dilution.

Antigen Retrieval: Trilogy™ (EDTA based, pH 8.0) buffer, 15min



For full product information, images and publications, please visit our website.

Date 2024 / 04 / 29 Page 2 of 2