

# p53 antibody

## Cat. No. GTX102965

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, ICC/IF, IHC-P, IP, ChIP assay
Reactivity	Human, Zebrafish



## Applications

## **Application Note**

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500-1:3000
ICC/IF	1:100-1:1000
IHC-P	1:100-1:1000
IP	1:100-1:500
ChIP assay	Assay dependent
Not tested in other applications.	

Calculated MW 44 kDa. (Note)

Form Liquid  Buffer PBS, 20% Glycerol	
Buffer PBS, 20% Glycerol	
Preservative 0.025% ProClin 300	
Storage Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4' long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.	°C. For
<b>Concentration</b> 1.28 mg/ml (Please refer to the vial label for the specific concentration.)	
Immunogen Recombinant protein encompassing a sequence within the center region of human p53. The exact sequence is proprie	etary.
Purification Purified by antigen-affinity chromatography.	
Conjugation Unconjugated	



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 11 / 02 Page 1 of 2



For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

#### Note


Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

#### DATA IMAGES



#### GTX102965 WB Image

Non-transfected (–) and transfected (+) 293T whole cell extracts (30  $\mu$ g) were separated by 10% SDS-PAGE, and the membrane was blotted with p53 antibody (GTX102965) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



#### GTX102965 ICC/IF Image


p53 antibody detects p53 protein at nucleus by immunofluorescent analysis.

Samples: HCT 116 cells mock (left) and treated with 30  $\mu$ M Cisplatin for 24 hrs (right) were fixed in 4% paraformaldehyde at RT for 15 min.

Green: p53 protein stained by p53 antibody (GTX102965) diluted at 1:500.

Blue: Hoechst 33342 staining.

Scale bar =  $10 \mu m$ .




### GTX102965 IHC-P Image

p53 antibody detects p53 protein at nucleus by immunohistochemical analysis.

Sample: Paraffin-embedded human breast carcinoma.

p53 stained by p53 antibody (GTX102965) diluted at 1:500.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min



### GTX102965 WB Image

Sample (30 µg of whole cell lysate)

A: HCT116 cells with mock treatment for 24 hr

B: HCT116 cells with 30  $\mu M$  cisplatin treatment for 24 hr

10% SDS PAGE

GTX102965 diluted at 1:1000

The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 11 / 02 Page 2 of 2