

# CDK4 antibody

# Cat. No. GTX102993

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, ICC/IF, IHC-P
Reactivity	Human, Mouse, Rat, Bovine



# Applications

## **Application Note**

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500-1:3000
ICC/IF	1:100-1:1000
IHC-P	1:100-1:1000
Not tested in other applications.	

Calculated MW 34 kDa. (Note)

Properties		
Form	Liquid	
Buffer	PBS, 20% Glycerol	
Preservative	0.025% ProClin 300	
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.	
Concentration	0.82 mg/ml (Please refer to the vial label for the specific concentration.)	
Immunogen	Recombinant protein encompassing a sequence within the center region of human CDK4. The exact sequence is proprietary.	
Purification	Purified by antigen-affinity chromatography.	
Conjugation	Unconjugated	
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.	
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.	




For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 11 / 02 Page 1 of 2



## DATA IMAGES



## GTX102993 ICC/IF Image


CDK4 antibody detects CDK4 protein at nucleus by immunofluorescent analysis.

Sample: MCF7 cells were fixed in 4% paraformaldehyde at RT for 15 min.

Green: CDK4 protein stained by CDK4 antibody (GTX102993) diluted at 1:1000.


Red: p21 Cip1, a nucleus marker, stained by p21 Cip1 antibody [GT1032] (GTX629543) diluted at 1:500.

Blue: Hoechst 33342 staining.




#### GTX102993 WB Image

Whole cell extract (30  $\mu$ g) was separated by 12% SDS-PAGE, and the membrane was blotted with CDK4 antibody (GTX102993) diluted at 1:500. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



#### GTX102993 WB Image

Whole cell extract (30  $\mu$ g) was separated by 12% SDS-PAGE, and the membrane was blotted with CDK4 antibody (GTX102993) diluted at 1:500. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



#### GTX102993 WB Image

Non-transfected (–) and transfected (+) 293T whole cell extracts (30  $\mu$ g) were separated by 12% SDS-PAGE, and the membrane was blotted with CDK4 antibody (GTX102993) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 11 / 02 Page 2 of 2