

UAP1 antibody [N1C3]

Cat. No. GTX103592

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, ICC/IF, IHC-P
Reactivity	Human

References (2)

Package

100 µl, 25 µl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500-1:3000
ICC/IF	1:100-1:1000
IHC-P	1:100-1:1000

Not tested in other applications.

Calculated MW 59 kDa. ([Note](#))

Properties


Form	Liquid
Buffer	0.1M Tris, 0.1M Glycine, 10% Glycerol
Preservative	0.01% Thimerosal
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant protein encompassing a sequence within the center region of human UAP1. The exact sequence is proprietary.
Purification	Purified by antigen-affinity chromatography.
Conjugation	Unconjugated

Note

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

For full product information, images and publications, please visit our [website](#).

DATA IMAGES

GTx103592 ICC/IF Image


UAP1 antibody [N1C3] detects UAP1 protein at cytoplasm by immunofluorescent analysis.

Sample: HeLa cells were fixed in 4% paraformaldehyde at RT for 15 min.

Green: UAP1 protein stained by UAP1 antibody [N1C3] (GTx103592) diluted at 1:200.


Red: alpha Tubulin, a cytoskeleton marker, stained by alpha Tubulin antibody [B-5-1-2] (GTx11304) diluted at 1:10000.

Blue: Hoechst 33342 staining.


GTx103592 IHC-P Image

Immunohistochemical analysis of paraaffin-embedded ES-2 xenograft , using UAP1(GTx103592) antibody at 1:500 dilution.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min



GTx103592 WB Image

Sample (30 ug of whole cell lysate)

A: H1299

7.5% SDS PAGE

GTx103592 diluted at 1:3000


GTx103592 ICC/IF Image

Immunofluorescence analysis of paraformaldehyde-fixed A549, using UAP1(GTx103592) antibody at 1:200 dilution.



For full product information, images and publications, please visit our [website](http://www.genetex.com).