

# Spi-C antibody

## Cat. No. GTX104182

| Host        | Rabbit     |
|-------------|------------|
| Clonality   | Polyclonal |
| Isotype     | IgG        |
| Application | WB         |
| Reactivity  | Human      |

Package  $100~\mu l, 25~\mu l$ 

## APPLICATION

### **Application Note**

\*Optimal dilutions/concentrations should be determined by the researcher.

| Suggested dilution                | Recommended dilution |
|-----------------------------------|----------------------|
| WB                                | 1:500-1:3000         |
| Not tested in other applications. |                      |

**Calculated MW** 29 kDa. ( <u>Note</u> )


| PROPERTIES    |  |
|---------------|--|
| Form          | Liquid   |
| Buffer        | PBS, 20% Glycerol  |
| Preservative  | 0.01% Thimerosal   |
| Storage       | Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles. |
| Concentration | 1 mg/ml (Please refer to the vial label for the specific concentration.)   |
| Immunogen     | Recombinant protein encompassing a sequence within the center region of human Spi-C. The exact sequence is proprietary.  |
| Purification  | Purified by antigen-affinity chromatography.   |
| Conjugation   | Unconjugated   |
| Note          | For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.  |
|               | Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.  |



For full product information, images and publications, please visit our website.

Date 2024 / 05 / 04 Page 1 of 2

### DATA IMAGES



#### GTX104182 WB Image

Non-transfected (–) and transfected (+) 293T whole cell extracts (30  $\mu$ g) were separated by 12% SDS-PAGE, and the membrane was blotted with Spi-C antibody (GTX104182) diluted at 1:500. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



For full product information, images and publications, please visit our <u>website</u>.

Date 2024 / 05 / 04 Page 2 of 2