

ZEB1 antibody [N2C1], Internal

Cat. No. GTX105278

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, ICC/IF, IP, ChIP assay
Reactivity	Human, Mouse, Rat



Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500-1:3000
ICC/IF	1:100-1:1000
IP	1:100-1:500
ChIP assay	Assay dependent

Not tested in other applications.

Calculated MW 124 kDa. (Note)

Properties	
Form	Liquid
Buffer	PBS, 1% BSA, 20% Glycerol
Preservative	0.025% ProClin 300
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.13 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant protein encompassing a sequence within the center region of human ZEB1. The exact sequence is proprietary.
Purification	Purified by antigen-affinity chromatography.
Conjugation	Unconjugated
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.




For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 12 / 06 Page 1 of 2




DATA IMAGES




GTX105278 IP Image

ZEB1 antibody immunoprecipitates ZEB1 protein in IP experiments. IP Sample: 293T whole cell lysate/extract A. 40 μ g 293T whole cell lysate/extract B. Control with 2 μ g of preimmune rabbit lgG C. Immunoprecipitation of ZEB1 protein by 2 μ g of ZEB1 antibody (GTX105278) 7.5% SDS-PAGE The immunoprecipitated ZEB1 protein was detected by ZEB1 antibody (GTX105278) diluted at 1:1000. EasyBlot anti-rabbit lgG (GTX221666-01) was used as a secondary reagent.



GTX105278 WB Image


Non-transfected (–) and transfected (+) HeLa whole cell extracts (30 μ g) were separated by 5% SDS-PAGE, and the membrane was blotted with ZEB1 antibody [N2C1], Internal (GTX105278) diluted at 1:500. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody, and the signal was developed with Trident ECL plus-Enhanced.



GTX105278 WB Image

Various whole cell extracts (30 μ g) were separated by 5% SDS-PAGE, and the membranes were blotted with ZEB1 antibody [N2C1], Internal (GTX105278) diluted at 1:1000 and competitor's antibody diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.

*The competitor is not affiliated with GeneTex and does not endorse this product.



GTX105278 ChIP assay Image

Cross-linked ChIP was performed with HeLa chromatin extract and 5 μ g of either control rabbit IgG or anti-ZEB1 antibody. The precipitated DNA was detected by PCR with primer set targeting to LAMC2 promotor. ChIP experiment and primer designs are based on <u>Proc Natl Acad Sci U S A. 2011 Nov 29;108(48):19204-9</u>.



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 12 / 06 Page 2 of 2