

Activin Receptor Type IIB antibody

Cat. No. GTX10596

| Host | Goat |
|--------------|----------------|
| Clonality | Polyclonal |
| Isotype | IgG |
| Applications | WB, IHC-P, FCM |
| Reactivity | Human |

Package $25 \, \mu g$

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

| Suggested dilution | Recommended dilution |
|-----------------------------------|-------------------------------------|
| WB | 0.1-0.2 μg/ml |
| IHC-P | 5 μg/ml |
| FCM | 10 μl/1-2.5 x 10 ⁵ cells |
| Not tested in other applications. | |

Calculated MW 58 kDa. (<u>Note</u>)

| Properties | |
|---------------|--|
| Form | Liquid |
| Buffer | PBS |
| Preservative | 0.02% Sodium azide |
| Storage | Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles. |
| Concentration | Batch dependent (Please refer to the vial label for the specific concentration.) |
| Immunogen | purified, NSO-derived, recombinant human activin receptor IIB extracellular domain. |
| Purification | Purified by affinity chromatography |
| Conjugation | Unconjugated |
| Note | For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption. |
| | Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product. |



For full product information, images and publications, please visit our website.

Date 2025 / 12 / 12 Page 1 of 1