

c-Met antibody

Cat. No. GTX10728

Host	Goat
Clonality	Polyclonal
Isotype	IgG
Applications	WB, FCM, IHC, Neutralizing/Inhibition
Reactivity	Human

Package
25 µg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	0.1-0.2 µg/ml
FCM	Assay dependent
IHC	5-15 µg/ml
Neutralizing/Inhibition	0.5-2 µg/ml

Not tested in other applications.

Calculated MW 156 kDa. ([Note](#))

Properties

Form	Liquid
Buffer	PBS carbohydrates
Preservative	No preservatives
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	Batch dependent (Please refer to the vial label for the specific concentration.)
Immunogen	purified recombinant human hepatocyte growth factor receptor extracellular domain expressed in mouse NSO cells.
Purification	Purified by affinity chromatography
Conjugation	Unconjugated

Note

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).



For full product information, images and publications, please visit our [website](#).